

Genetic Analysis of Human Immunodeficiency Virus Type 1 Strains in Kenya: A Comparison Using Phylogenetic Analysis and a Combinatorial Melting Assay

KENNETH E. ROBBINS,¹ LEONDIOS G. KOSTRIKIS,² TERESA M. BROWN,¹ OMU ANZALA,³
SUNNY SHIN,² FRANCIS A. PLUMMER,³ and MARCIA L. KALISH¹

ABSTRACT

We surveyed human immunodeficiency virus (HIV) subtype distribution from peripheral blood mononuclear cells (PBMCs) collected in 1995 from 24 HIV-1-infected Kenyan residents (specimens from predominantly male truck drivers and female sex workers near Mombasa and Nairobi). Processed lysates from the PBMC samples were used for *env* amplification, directly sequenced, and analyzed by phylogenetic analysis. Envelope amplification products were also used for analysis in a polymerase chain reaction (PCR)-based assay, called the combinatorial melting assay (COMA). Results of the two tests were compared for assignment of subtype for this Kenyan cohort. The COMA, a PCR capture technique with colorimetric signal detection, was used with HIV reference subtype strains as well as regional (East Africa) HIV strains for subtype identification. Performance of the COMA was at 100% concordance (24 of 24) as compared with DNA sequencing analysis. Phylogenetic analysis showed 17 isolates to be subtype A, 3 subtype D, and 4 subtype C viruses. This may represent an increase in subtype C presence in Kenya compared with previously documented reports. The COMA can offer advantages for rapid HIV-1 subtype screening of large populations, with the use of previously identified regional strains to enhance the identification of local strains. When more detailed genetic information is desired, DNA sequencing and analysis may be required.

INTRODUCTION

THE HUMAN IMMUNODEFICIENCY VIRUS TYPE 1 (HIV-1) global epidemic is currently thought to be composed of at least 10 phylogenetically distinct group M subtypes, designated A through J, and group O viruses.¹⁻⁴ Worldwide, and especially in African countries, where the epidemic is oldest, multiple subtypes predominate.^{5,6} The surveillance and documentation of strains, both established as well as newly disseminated in countries, are important for epidemiological tracking, diagnostic testing methodologies, protection of the blood supply, vaccine development, and field trials.⁷ Several techniques have been used to determine subtypes of HIV-1: (1) nucleotide sequencing,^{3,4} (2) the enzyme immunoassay utilizing peptides from the variable 3 region (V3) of the gp120 region from HIV-1,⁸⁻¹⁰ (3)

the heteroduplex mobility assay (HMA),^{11,12} and (4) subtype classification by polymerase chain reaction (PCR).¹³ While all of these methods have particular advantages and disadvantages, nucleotide sequencing is the most definitive and informative. However, compared with sequencing, the HMA has been demonstrated to be reliable for subtyping.¹⁴ The most frequently sequenced portion of the HIV-1 genome is the V3 and flanking regions of gp120 in the *env* gene.⁴ This is due to the biological importance of this area¹⁵⁻¹⁹ as well as to the large numbers of strains sequenced for this region and available for phylogenetic comparison in the Los Alamos National Laboratory HIV database.⁴ Sequence analysis of this gene region has been shown to provide accurate phylogenetic relationships, using most of the commonly available methods for inferring phylogenetic relationships.^{20,21} To determine the subtype distribu-

¹Division of AIDS, Sexually Transmitted Diseases, and Tuberculosis Laboratory Research, National Center for Infectious Diseases, Centers for Disease Control and Prevention, Public Health Service, U.S. Department of Health and Human Services, Atlanta, Georgia 30333.

²Aaron Diamond AIDS Research Center, Rockefeller University, New York, New York 10016.

³Department of Medical Microbiology, University of Manitoba, Winnipeg, Manitoba R3E 0W3, Canada.

Note: Use of trade names and commercial sources is for identification only and does not imply endorsement by the Public Health Service or the U.S. Department of Health and Human Services.

tion of HIV strains in truck drivers and sex workers near Mombasa and Nairobi, Kenya, and to better understand their contribution to the dissemination and mixing of subtypes, specimens from HIV-infected persons were obtained and sequenced directly from PCR fragments from the envelope gene. Phylogenetic analysis of the sequences was compared with the results produced by a new combinatorial melting assay (COMA),²² utilized here for HIV-1 subtype identification. The COMA subtypes an amplification product that encompasses the *env* region we sequenced, plus approximately 100 bp upstream and 100 bp downstream. The utility and performance characteristics of the two techniques are discussed.

MATERIALS AND METHODS

Specimens and isolation of viral DNA

Twenty-four HIV-1 specimens were obtained in 1995 from participants in a Kenyan sexual transmission cohort study, near Mombasa and Nairobi. The individuals were predominantly male truck drivers and female sex workers. The peripheral blood mononuclear cell (PBMC) processing and PCR amplification of the DNA lysates were as previously described,²³ except that the primary amplification primers for DNA sequencing were MK369 (5'-TGGAGCCAGTAGATCCTAGACTAGAGCCCT) and MK616 (5'-AATGGTGAGTATCCCTGCCTAAC-TCTATT) and the nested amplification primers for DNA sequencing were MK650 (5'-AATGTCAGCACAGTACAA-TGTACAC) and MK601 (5'-TTCTCCAATTGTCCTCATA-TCTCCTCCTCCA). The primary amplification primers utilized for the COMA PCR product were either MK369/MK616, MK603 (5'-CAGAAAATGGTGGGTACAGTCTATTATGGGGTACCT) and MK602 (5'-GCCCATAGTGCTTCCTGCTGCTCCCAAGAACC), or ED5/ED12.¹¹ Nested amplification primers^{2,22} for the COMA PCR products were LK1/LK2.²²

DNA sequencing

PCR-amplified products were purified by the Qiagen PCR purification kit (Qiagen, Chatsworth, CA). Purified DNA was sequenced according to the ABI PRISM dye terminator cycle sequencing Ready Reaction kit (Applied Biosystems/Perkin-Elmer, Foster City, CA). Sequencing primers used were various combinations of the PCR primers (MK616, MK650) and an internal sequencing primer KR207 (5'-CTGTTAAATGGCAGTCTAGC).

Phylogenetic analysis

The sequences were manually aligned with the sequence editor function of the Genetic Data Environment (GDE) package²⁴ and positions where gaps occurred were removed from analysis, leaving 327 nucleotide sites. Percentage diversity was calculated with the *pdist* function of the Molecular Evolution Genetic Analysis (MEGA) program.²⁵ Phylogenetic analysis was accomplished with PHYLIP (Phylogeny Inference Package)²⁶ for both neighbor-joining and maximum likelihood trees. Pairwise evolutionary distances were estimated for the neigh-

bor-joining trees by using the maximum likelihood option as the evolutionary model in the *dnadist* program. Utilizing the *consense* program, a consensus tree was developed from the 500 bootstrapped, neighbor-joining trees.

Nucleotide sequence accession numbers

The GenBank accession numbers for the Kenyan sequences reported in this study are listed in Table 1.

COMA

The COMA was performed by two of us (L.K. and S.S.) as previously described,²² with modifications of the reference strains used. Briefly, antisense, single-stranded DNAs (ssDNAs) were generated for reference subtypes A, B, C, and D by asymmetrical PCR, with biotinylation of one of the primers facilitating capture to streptavidin-coated plates. Reference and regional strains used for the different subtypes (followed by each GenBank accession number) were as follows: A (92KEML1048:AF103913, 91KEMCH9727:AF103910, 91KEMCHI96:AF103911, and 92KEML639:AF103914), B (BR20:U08797 and TH14:U08801), C (SM145:L22946, DJ259:L22940, and MW6510:L15733), and D (91KEML22:AF10312, 89KEML203:AF103909, UG21:U27399, and UG46:U08809). Reference PCR products from the same subtype were combined to form reference pools and placed in rows of the microtiter plate. Complementary sense-stranded molecules from the Kenyan DNA were produced by asymmetrical PCR with digoxigenin labeling. These samples were placed in the microtiter plate columns, allowing heteroduplex formation between the Kenyan DNA and reference subtype DNA. The subtype was determined by a colorimetric immunoassay using alkaline phosphatase-conjugated antibody against digoxigenin. The correct subtype was determined by the comparative absorbance measurement, rounded to the first decimal place, being higher than any other by a factor of at least two.

RESULTS

PCR amplification and nucleotide analysis

PCR amplification of the C2V3 region of the *env* gene was performed with two sets of primers in a nested fashion in two separate PCRs. Nucleotide sequences were obtained directly from the PCR product and aligned for homologous positions, and a neighbor-joining phylogenetic tree was constructed (Fig. 1). Seventeen of the Kenyan sequences clustered within the subtype A group, 4 with subtype C, and 3 with subtype D. The same subtype classifications were obtained with a tree constructed with the maximum likelihood program (data not shown). Two pairs of sequences (KEM11/KEQRS1, KETK5/KEM14) clustered together with bootstrap support greater than 70% (Fig. 1) as well as in the maximum likelihood tree. These sequences are from female sex workers and a female sex worker/male truck driver, respectively (see caption to Fig. 1). Whether these pairs represent epidemiologically linked infections is unknown, yet the sequences within each pair being derived from individuals residing in different cities (Mombasa and Nairobi) suggests some infection links between these commu-

TABLE 1. COMA VERSUS SEQUENCING SUBTYPING RESULTS OF THE STUDY SPECIMENS

Isolate	GenBank accession numbers	<i>Relative absorbance densities (490 nm) of DNA heteroduplexes formed by complementary ssDNA from each unknown env and four reference env repertoires</i>				Subtype determined by COMA	Subtype determined by DNA sequencing
		A	B	C	D		
KEMCH1	AF103916	1.00	0.12	0.10	0.13	A	A
KEMCH3	AF103917	1.00	0.09	0.13	0.11	A	A
KEMCH4	AF103918	1.00	0.13	0.13	0.13	A	A
KEML3	AF103921	1.00	0.08	0.09	0.10	A	A
KEML4	AF103922	1.00	0.08	0.08	0.08	A	A
KEML5	AF103923	1.00	0.17	0.17	0.21	A	A
KEML6	AF103924	1.00	0.20	0.13	0.48	A	A
KEQRS2	AF103926	1.00	0.17	0.22	0.24	A	A
KEQRS4	AF103928	1.00	0.11	0.10	0.15	A	A
KETDP1	AF103930	1.00	0.12	0.14	0.18	A	A
KETDP2	AF103931	1.00	0.11	0.13	0.13	A	A
KETK2	AF103934	1.00	0.07	0.11	0.10	A	A
KETK3	AF103935	1.00	0.11	0.12	0.12	A	A
KETK4	AF103936	1.00	0.10	0.16	0.14	A	A
KETK5	AF103937	1.00	0.07	0.17	0.08	A	A
KESHH1	AF103929	1.00	0.07	0.04	0.11	A	A
KEX1	AF103938	1.00	0.07	0.09	0.10	A	A
KEML1	AF103919	0.27	0.12	1.00	0.18	C	C
KEQRS1	AF103925	0.28	0.28	1.00	0.23	C	C
KETDP3	AF103932	0.19	0.26	1.00	0.18	C	C
KETK1	AF103933	0.51	0.07	1.00	0.24	C	C
KEMCH2	AF103916	0.26	0.46	0.11	1.00	D	D
KEML2	AF103920	0.09	0.08	0.11	1.00	D	D
KEQRS3	AF103927	0.11	0.08	0.12	1.00	D	D

nities. Subtypes A and C are represented in these two pairs, and all three subtypes found in this study have previously been reported in Kenya.^{6,27,28}

The percentage diversity of the nucleotide sequences was calculated for the Kenyan taxa in Fig. 1. The mean genetic distance among Kenyan subtype A sequences is 13.2%, while the subtype C distance is 11.2% and the subtype D distance is 10.7%. An alignment of the predicted amino acid sequences for the 24 Kenyan sequences is depicted in Fig. 2. All are compared with subtype-specific consensus sequences from the Los Alamos HIV database.⁴

The GPGQ tetrameric tip of the V3 loop was predominant in all three subtypes, with the exception of two type As (KESHH1, KEX1) and one type D (KEQRS3) having GPGR. Positively charged amino acids at positions 11 and 25 of the V3 loop, generally believed to be associated with a syncytium-inducing phenotype,²⁹ were absent in all of the Kenya sequences. This would be expected for subtype A strains, but positively charged amino acids at one or two of these positions usually make up the majority of published African type D strains.^{6,14,30} One explanation could be that these Kenyan strains may be from individuals in an early stage of infection,^{31,32} although clinical data are not available.

COMA and comparison with sequencing

PCR amplification of the C2V4 region of the *env* gene was performed with one set of primers and subsequently sent to two

of us (L.K. and S.S.)²² for secondary amplification and COMA testing (see Materials and Methods). Table 1 lists the COMA results by relative absorbance densities of the 24 Kenyan specimens in comparison with the subtype determined by sequencing and phylogenetic analysis. All specimens are in agreement between the two methods, corresponding to a 100% concordance (24 of 24).

DISCUSSION

COMA versus sequencing analysis for HIV-1 subtype identification

Along with investigating the distribution of strains among the highly mobile cohort of truck drivers and sex workers, we compared the accuracy of the COMA with DNA sequencing and subsequent phylogenetic analysis and adapted the COMA in a region-specific fashion for subtyping HIV strains from Kenya. Phylogenetic analysis was performed on a genomic region within and approximately 200 bp shorter than the PCR fragment used in the COMA. The extensive use of this region by us and others for accurate subtyping allowed the utility of sequencing this shorter fragment for comparisons with the COMA results. The complete concordance for this Kenyan HIV-1 sampling between the two methodologies (100%, 24 of 24) demonstrated that the two *env* regions could be compared reliably by these methods (although the sample size is relatively




FIG. 1. Phylogenetic analysis of 24 HIV-1 DNA specimens obtained from individuals in Kenya. Analysis is based on 327 alignable positions (positions in alignment containing one or more gaps are excluded from analysis) in the C2V3 region of the HIV-1 *env* gene with the simian immunodeficiency virus SIV_{cpzgab} as outgroup. Los Alamos HIV Database subtype sequences from A, B, C, D, E, F, G, and H are included to facilitate subtype clustering of the Kenyan sequences. The Kenyan sequences determined in this study (boldface with a KE prefix, isolation date of 1995) are designated as follows: MCH, female sex workers—region unknown; ML, female sex workers from Nairobi; QRS, female sex workers from Mombasa; TDP, male truck drivers near Nairobi; TK, male truck drivers near Mombasa; SHH, male sexually transmitted disease patient; X, unknown. For clarity, those originating from Mombasa have an -m suffix and those from Nairobi have an -n suffix. The numbers at the branch points indicate the percentage of times the branch was preserved in the 500 bootstrap replications. Only bootstrap proportions >70% are shown. The large letters indicate A–H subtypes.




FIG. 2. Amino acid alignment of the *env* C2V3 regions of the Kenyan strains compared with subtype A, C, and D consensus sequences.²⁰ Kenyan descriptive names are as listed in the caption to Fig. 1. A dash (–) indicates identity with the A, C, or D consensus sequence and a period (.) depicts a deletion.

small) and supports the use of the COMA to subtype previously unknown samples. A key feature of the COMA is its flexible use of multiple strains from each subtype for detection of the unknown strain. Utilization of strains found in the geographic region of interest as reference strains is a customization of the more general COMA, which is run with worldwide reference strains for subtyping.²² Specimens from subtypes A and D, which have been reported to predominate in Kenya and Uganda, were used in the cocktail of reference strains. The Kenyan reference strains were from female sex workers, isolated from 1989 to 1992. Further refinement of signal-to-noise ratios for the COMA may be assisted by addition of more recently subtyped specimens to the respective subtype reference pool. As we demonstrated here, a population-based study could be performed using the COMA technology for initial subtype identification, without DNA sequencing. Ambiguous or cross-reacting specimens (i.e., subtype recombinants or dual infections) could then be sequenced for further identification, if needed.

In addition, the COMA can be used to systematically monitor change in subtype distribution over time in large population-based surveys. Immunology-based assays such as peptide-binding enzyme immunoassays (EIAs) have been effective where the epidemic was relatively new and, consequently, the diversity within subtypes was low, e.g., in Thailand.⁹ In regions where the epidemic is older and people have been infected for longer periods, the within person distribution of quasispecies is more complex, and the host immune system has responded to many antigenically distinct variants. These samples typically have a large degree of cross-reactivity in serological peptide-based assays. Therefore, the HMA^{11,14} and COMA techniques, as demonstrated previously²² and in this study, are more reliable for subtype assignment. Subtype analysis with the HMA typically uses two reference strains per subtype, while the microtiter format of the COMA allows many reference strains to be used with no increase in the number of wells. Sequencing

and subsequent phylogenetic analysis, although more time consuming, is generally considered the “gold standard.” The COMA has the advantages of using a format typical of that widely used in EIA technology, requiring little equipment beyond a thermocycler for PCR, and being easy to interpret. The HMA could be used as a complement to initial COMA screening by identifying major and minor quasispecies of samples that yield indeterminate COMA results. Alternatively, or as the next step, the PCR-amplified specimen could be sequenced for definitive typing, mutation analysis, amino acid sequence prediction, and identification of recombinant viruses.

Kenyan HIV-1 subtypes

HIV-1 subtyping has been an important part of the surveillance and characterization of HIV strains for designing vaccine strategies prior to conducting vaccine field trials. Many African countries harbor a large variety of subtypes represented in the global epidemic. In particular, the Central and East African countries, which includes Kenya, compose that region of the continent where most of the subtypes have been reported.^{14,33} Our interest in tracking the spread of HIV-1 subtypes in this area led us to obtain and analyze 24 specimens from a Kenyan sex worker/truck driver cohort. The finding that most (17 of 24) of these strains grouped with subtype A and a minority with subtype D (3 of 24) is similar to previously published reports of Kenyan sequences⁶ and is also consistent with the rapid spread of subtype A in Central Africa.¹⁴ Subtype C (4 of 24 in this study) is also well established in several southern African countries,^{7,34,35} as well as in India.³⁶ The spread of this subtype to other countries includes China,³⁷ Malaysia,³⁸ Russia,³⁹ England,⁴⁰ and the United States (P. Sullivan, Centers for Disease Control [CDC, Atlanta, GA], unpublished data, 1998). The United Nations Program on AIDS (UNAIDS, Geneva, Switzerland) has identified subtypes A and C as the most prevalent

group M strains worldwide and has organized collaborative international studies for better sampling and characterization of these subtypes.⁴¹ The identification of the C subtype in these Kenyan residents is further documentation of its continuing spread. Previous Kenyan HIV subtype sampling studies^{6,27,28} for specimens collected from 1991 to 1994 had demonstrated only one subtype C strain. Although the sampling numbers are relatively small, the increased number of subtype C virus strains identified here could indicate an increasing prevalence of subtype C HIV strains in Kenya. It at least establishes that the initial report²⁸ of a subtype C virus in Kenya was not spurious and that subtype C viruses are being transmitted in this region.

The worldwide trend toward an increase in multiple subtypes in regions where only a few previously predominated is typically associated with increased travel and demographic shifts.⁴² Truck drivers involved in this study transport goods along the trans-African highway, beginning in Mombasa (a major regional seaport), with final destinations in Uganda, Tanzania, Rwanda, Burundi, eastern Zaire, and southern Sudan.⁴³ The drivers are on the road for many weeks or months and have frequent contact with female sex workers along the route. Sex workers are also mobile along the truck route, sometimes accompanying drivers. Both drivers and sex workers have high rates of HIV-1 infection. Thus, these individuals may be exposed to HIV-1 subtypes from a wide geographic area and could provide a means of introducing new HIV-1 subtypes (subtype C in this case) into new geographic areas. The observation that two (A and C) of the three subtypes found here have phylogenetically linked sequences from residents of different Kenyan cities may further indicate the effectiveness of spread by the high-risk behavior within these risk groups.

REFERENCES

- De Leys R, Van der Borgh B, Vanden Haesevelde M, Heyndrickx L, VanGeel A, Wauters C, Bernaerts R, Saman E, Nijs P, Willems B, Taelman H, van der Goen G, Piot P, Tersmette T, Huisman JG, and Van Heuverswyn H: Isolation and partial characterization of an unusual human immunodeficiency retrovirus from 2 persons of West-Central African origin. *J Virol* 1990;64:1207-1216.
- Kostrikis LG, Bagdades E, Cao Y, Zhang L, Dimitriou D, and Ho DD: Genetic analysis of human immunodeficiency virus type 1 strains from patients in Cyprus: Identification of a new subtype designated subtype I. *J Virol* 1995;69:6122-6130.
- Louwagie J, McCutchan F, Peeters M, Brennan T, Sanders-Buell E, Eddy G, van der Groen G, Franssen K, Gershy-Damet G, Deleys R, and Burke D: Phylogenetic analysis of *gag* genes from 70 international HIV-1 isolates provides evidence for multiple genotypes. *AIDS* 1993;7:769-780.
- Myers G, Korber B, Hahn BH, Jeang K-T, Mellors JW, McCutchan FE, Henderson LE, and Pavlakis GN: *Human Retroviruses and AIDS: A Compilation and Analysis of Nucleic Acid and Amino Acid Sequences*. Los Alamos National Laboratory, Los Alamos, New Mexico, 1995.
- Bruce C, Clegg C, Featherstone A, Smith J, Biryahawaho B, Downing R, and Oram J: Presence of multiple genetic subtypes of human immunodeficiency virus type 1 proviruses in Uganda. *AIDS Res Hum Retroviruses* 1994;10:1543-1550.
- Janssens W, Heyndrickx L, Franssen K, Temmerman M, Leonaers A, Ivens T, Motte J, Piot P, and Van Der Groen G: Genetic variability of HIV type 1 in Kenya. *AIDS Res Hum Retroviruses* 1994;10:1577-1579.
- Hu DJ, Dondero TJ, Rayfield MA, George JR, Schochetman G, Jaffe HW, Luo C-C, Kalish ML, Weniger BG, Pau C-P, Schable CA, and Curran JW: The emerging genetic diversity of HIV. *JAMA* 1996;275:210-216.
- Cheingsong-Popov R, Lister S, Callow D, Kaleebu P, Weber S, and the WHO Network for HIV Isolation and Characterization: Serotyping HIV type 1 by antibody binding to the V3 loop: Relationship to viral genotype. *AIDS Res Hum Retroviruses* 1994;10:1379-1386.
- Pau C-P, Lee-Thomas S, Auwanit W, George JR, Ou C-Y, Parekh BS, Granade TS, Holloman DL, Phillips S, Schochetman G, Young NL, Takebe Y, Gayle HD, and Weniger BG: Highly specific V3 peptide enzyme immunoassay for serotyping HIV-1 specimens from Thailand. *AIDS* 1993;7:337-340.
- Wasi C, Herring B, Raktham S, Vanichseni S, Mastro TD, Young NL, Rübbsamen-Waigmann H, von Briesen H, Kalish ML, Luo C-C, Pau C-P, Baldwin A, Mullins JI, Delwart EL, Esparza J, Heyward WL, and Osmanov S: Determination of HIV-1 subtypes in injecting drug users in Bangkok, Thailand, using peptide binding enzyme immunoassay and the heteroduplex mobility assay: evidence of increasing infection with HIV-1 subtype E. *AIDS* 1995;9:843-849.
- Delwart EL, Shpaer EG, Louwagie J, McCutchen FE, Grez M, Rübbsamen-Waigmann H, and Mullins JI: Genetic relationships determined by a DNA heteroduplex mobility assay: Analysis of HIV-1 *env* genes. *Science* 1993;262:1257-1261.
- Osmanov S, Belsey L, Heyward W, Esparza J, Bradac B, Galva-Castro P, Van de Peere P, Karita E, Sempala S, Tugume B, Biryahawaho B, Wasi C, Rübbsamen-Waigmann H, von Briesen H, Esser U, Grez H, Holmes H, McCutchan FE, Louwagie J, Hegerich P, Lopez-Galindez C, Mullins JI, Delwart EL, Bachman MH, and Goudsmit J: HIV variation in WHO-sponsored vaccine evaluation sites: Genetic screening, sequence analyses and preliminary biological characterization of representative strains. *AIDS Res Hum Retroviruses* 1994;10:1327-1343.
- McCutchan FE, Sanders-Buell E, Oster CW, Redfield RR, Hira SK, Perine PL, Ungar BL, and Burke DS: Genetic comparison of human immunodeficiency virus (HIV-1) isolates by polymerase chain reaction. *J Acquir Immune Defic Syndr* 1991;4:1241-1250.
- WHO Network for HIV Isolation and Characterization: HIV type 1 variation in World Health Organization-sponsored vaccine evaluation sites: Genetic screening, sequence analysis, and preliminary biological characterization of selected viral strains. *AIDS Res Hum Retroviruses* 1994;10:1327-1343.
- DeJong JJ, Keulen W, Klaver B, Krone W, Tersmette M, and deRonde A: Human immunodeficiency virus type 1 clones chimeric for the envelope V3 domain differ in syncytium formation and replication capacity. *J Virol* 1992;66:757-765.
- Goudsmit J, Back NK, and Nara PL: Genomic diversity and antigenic variation of HIV-1: Links between pathogenesis, epidemiology and vaccine development. *FASEB J* 1991;5:2427-2436.
- Hwang SS, Boyle TJ, Lyerly HK, and Cullen BR: Identification of the envelope V3 loop as the primary determinant of cell tropism in HIV-1. *Science* 1991;253:71-74.
- Ivanhoff LA, Dubay JW, Morris JF, Roberts SJ, Gutshall L, Sternberg EJ, Hunter, Matthews TJ, and Petteway SR Jr: V3 loop region of the HIV-1 gp 120 envelope protein is essential for virus infectivity. *Virology* 1992;187:423-432.
- Javaherian K, Langlois AJ, LaRosa GJ, Profy AT, Bolognesi DP, Herlihy WC, Putney SD, and Matthews TJ: Broadly neutralizing antibodies elicited by the hypervariable neutralizing determinant in HIV-1. *Science* 1990;250:1590-1593.
- Hillis DM, Huelsenbeck JP, and Cunningham CW: Application and accuracy of molecular phylogenies. *Science* 1994;264:671-677.

21. Leitner T, Kescanilla D, Franzen C, Uhlen M, and Albert J: Accurate reconstruction of a known HIV-1 transmission history by phylogenetic tree analysis. *Proc Natl Acad Sci USA* 1996;93:19864-10869.
22. Kostrikis LG, Shin S, and Ho DD: Genotyping HIV-1 and HCV strains by a combinatorial DNA melting assay (COMA). *Mol Med* 1998;4:443-453.
23. Potts KE, Kalish ML, Bandea C, Orloff G, St Louis M, Brown C, Malanda N, Kavuka M, Schochetman G, Ou C-Y, and Heyward WL: Genetic diversity of human immunodeficiency virus type 1 (HIV-1) strains in Kinshasa, Zaire. *AIDS Res Hum Retroviruses* 1993;9:613-618.
24. Smith S: *GDE: Genetic Data Environment*, version 2.2. Millipore Imaging Systems, Ann Arbor, Michigan, 1992.
25. Kumar S, Tamura K, and Nei M: *MEGA: Molecular Evolutionary Genetic Analysis*, version 1.01. Pennsylvania State University, University Park, Pennsylvania, 1993.
26. Felsenstein J: *PHYLIP: Phylogeny Inference Package*, versions 3.5c and 4.0. University of Washington, Seattle, Washington, 1993.
27. Poss M, Gosink J, Thomas E, Kreiss JK, Ndinya-Achola J, Mandaliya K, Bwayo J, and Overbaugh J: Phylogenetic evaluation of Kenyan HIV type 1 isolates. *AIDS Res Hum Retroviruses* 1997;13:493-499.
28. Zachar V, Goustin AS, Zacharova V, Hager H, Koppelhus U, Womble DD, Liu X, Bamba C, Nyongo A, and Ebbesen P: Genetic polymorphism of envelope V3 region of HIV type 1 subtypes A, C, and D from Nairobi, Kenya. *AIDS Res Hum Retroviruses* 1996;12:75-78.
29. Milich L, Margolin B, and Swanstrom R: V3 Loop of the human immunodeficiency virus type 1 Env protein: Interpreting sequence variability. *J Virol* 1993;67:5623-5634.
30. Robbins KE, Bandea CI, Levin A, Goedert JJ, Blattner WA, Brubaker G, Brown TM, Schochetman G, Kalish ML, Shao J, and O'Brien TR: Genetic variability of human immunodeficiency virus type 1 in rural northwest Tanzania. *AIDS Res Hum Retroviruses* 1996;12:1389-1391.
31. Asjo BL, Morfeldt-Manson J, Biberfeld AG, Karlsson K, Lidman K, and Fenyö E-M: Replicative properties of human immunodeficiency virus from patients with varying severity of HIV infection. *Lancet* 1986;ii:660-662.
32. Tersmette M, De Goede R, Al BJM, Winkel I, Coutinho RA, Cuypers H, Huisman GJ, and Miedema F: Differential syncytium-inducing capacity of HIV isolates. Frequent detection of syncytium-inducing isolates in patients with AIDS and ARC. *J Virol* 1988;62:2026-2032.
33. Louwagie J, Janssens W, Mascola J, Heyndrickx L, Hegerich P, van der Groen G, McCutchan FE, and Burke DS: Genetic diversity of the envelope glycoprotein from human immunodeficiency virus type 1 isolates of African origin. *J Virol* 1995;69:263-271.
34. Murphy E, Korber B, Georges-Courbot MC, You B, Pinter A, Cook D, Kiény MP, Georges A, Mathiot C, and Barré-Sinoussi F: Diversity of the V3 region sequences of human immunodeficiency virus type 1 from the Central African Republic. *AIDS Res Hum Retroviruses* 1993;9:997-1006.
35. Orloff GM, Kalish ML, Chipangwi J, Potts KE, Ou C-Y, Schochetman G, Dallabetta G, Saah A, and Miotti PG: V3 Loops of HIV-1 specimens from pregnant women in Malawi uniformly lack a potential N-linked glycosylation site. *AIDS Res Hum Retroviruses* 1993;9:705-706.
36. Jameel S, Zafrullah M, Ahmad M, Kapoor GS, and Sehgal S: A genetic analysis of HIV-1 from Punjab, India reveals the presence of multiple variants. *AIDS* 1995;9:685-690.
37. Luo CC, Tian C, Hu DJ, Kai M, Dondero T, and Zheng X: HIV subtype in China. *Lancet* 1995;345:1051-1052.
38. Brown TM, Robbins KE, Sinniah M, Saraswathy TS, Lee V, Hooi LS, Vijayamalar B, Luo C-C, Ou C-Y, Rapier J, Schochetman G, and Kalish ML: HIV type 1 subtypes in Malaysia include B, C, and E. *AIDS Res Hum Retroviruses* 1996;12:1655-1657.
39. Lukashov VV, Cornelissen MTE, Goudsmit J, Papuashvilli MN, Rytik PG, Khaitov RM, Karamov EV, and de Wolf F: Simultaneous introduction of distinct HIV-1 subtypes into different risk groups in Russia, Byelorussia and Lithuania. *AIDS* 1995;9:435-439.
40. Arnold C, Barlow KL, Parry JV, and Clewey JP: At least five HIV-1 sequence subtypes (A, B, C, D, A/E) occur in England. *AIDS Res Hum Retroviruses* 1995;11:427-429.
41. European Commission (DG XII, INCO-DC) and the Joint United Nations Programme on HIV/AIDS: HIV-1 Subtypes: Implications for epidemiology, pathogenicity, vaccines and diagnostics. *AIDS* 1997;11:17-36. [Workshop report]
42. Muller O, Ungchusak K, Leng HB, Chung A, and Tadiar F: HIV and AIDS in southeast Asia. *Lancet* 1997;350:288.
43. Bwayo J, Plummer F, Omari M, Mutere A, Moses S, Ndinya-Achola J, Velentgas P, and Kreiss J: Human immunodeficiency virus infection in long-distance truck drivers in East Africa. *Arch Intern Med* 1994;154:1391-1396.

Address reprint requests to:

Kenneth E. Robbins

DASTLR, NCID

Centers for Disease Control and Prevention

1600 Clifton Rd. Mail Stop G-19

Atlanta, Georgia 30333

This article has been cited by:

1. Esmeralda AJM Soares, Ana MB Mart??nez, Thatiana M Souza, Andr?? FA Santos, Vanusa Da Hora, Jussara Silveira, Francisco I Bastos, Amilcar Tanuri, Marcelo A Soares. 2005. HIV-1 subtype C dissemination in southern Brazil. *AIDS* **19**:suppl 4, S81??S86. [[CrossRef](#)]
2. Marcelo Soares, Tulio de Oliveira, Rodrigo Brindeiro, Ricardo Diaz, Ester Sabino, Luis Brigido, Ivone Pires, Mariza Morgado, Maria Dantas, Draurio Barreira. 2003. *AIDS* **17**:1, 11. [[CrossRef](#)]
3. Dowling et al. 2003. Forty-one near full-length HIV-1 sequences from Kenya reveal an epidemic of subtype A and A-containing recombinants,. *AIDS* **17**:supplement 4, S125. [[CrossRef](#)]
4. Esmeralda A. J. M. Soares, Rodrigo P. Santos, Jos?? Augusto Pellegrini, Eduardo Sprinz, Amilcar Tanuri, Marcelo A. Soares. 2003. Epidemiologic and Molecular Characterization of Human Immunodeficiency Virus Type 1 in Southern Brazil. *JAIDS Journal of Acquired Immune Deficiency Syndromes* **34**:5, 520. [[CrossRef](#)]
5. William Dowling, Bohye Kim, Carl Mason, K. Wasunna, Uzma Alam, Lynne Elson, Deborah Birx, Merlin Robb, Francine McCutchan, Jean Carr. 2002. *AIDS* **16**:13, 1809. [[CrossRef](#)]
6. Jos?? Esparza, Donald Burke. 2001. Epidemiological considerations in planning HIV preventive vaccine trials. *AIDS* **15**:supplement 5, S49. [[CrossRef](#)]